

SUPER TRIPLE SYSTEMS AND APPLICATIONS TO PARA-STATISTICS AND YANG-BAXTER EQUATION

S. OKUBO

*Department of Physics and Astronomy, University of Rochester
Rochester, NY 14627, USA*

ABSTRACT

We introduce the notion of ortho-symplectic super triple system, and apply it to find solutions of super Yang-Baxter equation. Also, the para-statistics are formulated as a Lie-super triple system.

1. Triple Product

1.1 Quaternionic and Octonionic Triple Systems

Let V be a N -dimensional vector space, which we write hereafter as

$$N = \text{Dim } V \quad . \quad (1.1)$$

A triple product $w = [x, y, z]$ in V is an assignment of $w \in V$ for any three elements $x, y, z \in V$, which is linear in each variables x, y , and z . In other words, we may write

$$[x, y, z] ; V \otimes V \otimes V \rightarrow V \quad . \quad (1.2)$$

Perhaps the simplest non-trivial example is the case of $N = 4$, which can be constructed below. Let e_1, e_2, e_3 , and e_4 be a basis of V and assign $[e_j, e_k, e_\ell] \in V$ by

$$[e_j, e_k, e_\ell] = \sum_{m=1}^4 \epsilon_{jklm} e_m \quad (1.3)$$

for any $j, k, \ell = 1, 2, 3, 4$, where ϵ_{jklm} is the totally antisymmetric Levi-Civita symbol with $\epsilon_{1234} = 1$ in 4-dimensional space. We then extend the definition of $[x, y, z]$ for any $x, y, z \in V$ from Eq. (1.3) by the linearity. Further, we can introduce a bi-linear form $\langle x|y \rangle$ in V by

$$\langle e_j|e_k \rangle = \delta_{jk} \quad (j, k = 1, 2, 3, 4) \quad . \quad (1.4)$$

Then, $\langle x|y \rangle$ defines clearly a symmetric bi-linear non-degenerate form in V . The triple product $[x, y, z]$ with the bi-linear form $\langle x|y \rangle$ can be easily verified to satisfy the following properties:

$$(i) \quad \langle x|y \rangle = \langle y|x \rangle \text{ is non-degenerate,} \quad (1.5a)$$

$$(ii) \quad [x, y, z] \text{ is totally antisymmetric in } x, y, z, \quad (1.5b)$$

$$(iii) \quad \langle w|[x, y, z] \rangle \text{ is totally antisymmetric in } w, x, y, z, \quad (1.5c)$$

$$(iv) \quad \begin{aligned} & \langle [x, y, z][u, v, w] \rangle \\ &= \langle x|u \rangle \langle y|v \rangle \langle z|w \rangle + \langle x|v \rangle \langle y|w \rangle \langle z|u \rangle \\ &+ \langle x|w \rangle \langle y|u \rangle \langle z|v \rangle - \langle x|w \rangle \langle y|v \rangle \langle z|u \rangle \\ &- \langle x|v \rangle \langle y|u \rangle \langle z|w \rangle - \langle x|u \rangle \langle y|w \rangle \langle z|v \rangle, \end{aligned} \quad (1.5d)$$

$$(v) \quad \begin{aligned} & [u, v, [x, y, z]] \\ &= (\langle y|v \rangle \langle z|u \rangle - \langle y|u \rangle \langle z|v \rangle)x \\ &+ (\langle z|v \rangle \langle x|u \rangle - \langle z|u \rangle \langle x|v \rangle)y \\ &+ (\langle x|v \rangle \langle y|u \rangle - \langle x|u \rangle \langle y|v \rangle)z. \end{aligned} \quad (1.5e)$$

For examples, Eq. (1.5c) is an immediate consequence of

$$\langle e_m|[e_j, e_k, e_\ell] \rangle = \epsilon_{jk\ell m}$$

being totally antisymmetric in j, k, ℓ , and m , while Eqs. (1.5d) and (1.5e) is equivalent to the validity of the identity,

$$\begin{aligned} \sum_{m=1}^4 \epsilon_{jk\ell m} \epsilon_{abcm} &= \delta_{ja}(\delta_{kb}\delta_{\ell c} - \delta_{kc}\delta_{\ell b}) \\ &+ \delta_{jb}(\delta_{kc}\delta_{\ell a} - \delta_{ka}\delta_{\ell c}) + \delta_{jc}(\delta_{ka}\delta_{\ell b} - \delta_{kb}\delta_{\ell a}) \quad . \end{aligned}$$

Conversely, we can prove that any system satisfying Eqs. (1.5) is possible only for $\text{Dim } V = 4$, and moreover that we can find a basis e_1, e_2, e_3 and e_4 which satisfy Eqs. (1.3) and (1.4). We call the system to be a quaternionic triple (or ternary) system¹.

We can generalize Eqs. (1.5) by adding extra terms

$$\begin{aligned} & \beta \{ \langle x|u \rangle \langle y|[z, v, w] \rangle \\ &+ \langle y|u \rangle \langle z|[x, v, w] \rangle + \langle z|u \rangle \langle x|[y, v, w] \rangle \\ &+ \langle x|v \rangle \langle y|[z, w, u] \rangle + \langle y|v \rangle \langle z|[x, w, u] \rangle \\ &+ \langle z|v \rangle \langle x|[y, w, u] \rangle + \langle x|w \rangle \langle y|[z, u, v] \rangle \\ &+ \langle y|w \rangle \langle z|[x, u, v] \rangle + \langle z|w \rangle \langle x|[y, u, v] \rangle \} \quad , \end{aligned} \quad (1.5d)'$$

to the right side of Eq. (1.5d) and

$$\begin{aligned} & -\beta\{ < u|[v, y, z] > x + < u|[v, z, x] > y + < u|[v, x, y] > z\} \\ & -\beta\{ < x|v > [u, y, z] + < y|v > [u, z, x] + < z|v > [u, x, y] \\ & + < x|u > [v, z, y] + < y|u > [v, x, z] + < z|u > [v, y, x]\} \quad , \end{aligned} \quad (1.5e)'$$

to the right side of Eq. (1.5e) for a constant β , while other relations Eqs. (1.5a), (1.5b), and (1.5c) remain unchanged. We have shown elsewhere¹ that the new system is possible only for two cases of

$$(a) \quad N = 4 \quad , \quad \beta = 0 \quad (1.6a)$$

$$(b) \quad N = 8 \quad , \quad \beta = \pm 1 \quad . \quad (1.6b)$$

We call the new case of $N = \text{Dim } V = 8$ to be an octonionic triple system. Let us normalize β to be $\beta = -1$ for the octonionic triple system by changing the sign of $[x, y, z]$ if necessary. Let $e \in V$ be any fixed element satisfying

$$< e|e > = 1 \quad , \quad (1.7a)$$

and introduce a bi-linear product xy in V by

$$xy = [x, y, e] + < x|e > y + < y|e > x - < x|y > e \quad . \quad (1.7b)$$

Then, it is easy to show the validity of

$$xe = ex = x \quad , \quad (1.8a)$$

$$< xy|xy > = < x|x > < y|y > \quad (1.8b)$$

so that the bi-linear product xy defines the 8-dimensional octonion algebra². Conversely, we can determine the original triple product $[x, y, z]$ in terms of the bi-linear octonionic product, although we will not go into detail.

1.2 Orthogonal Triple System

Hereafter in this note, we assume that $< x|y >$ is a bi-linear non-degenerate form in V which is, however, not necessarily symmetric. Also, let $e_1, e_2, \dots, e_N$ be a basis of V and set

$$g_{jk} = < e_j|e_k > \quad . \quad (1.9)$$

Because of the non-degeneracy of $< x|y >$, g_{jk} possesses its inverse g^{jk} satisfying

$$\sum_{\ell=1}^N g_{j\ell} g^{\ell k} = \sum_{\ell=1}^N g^{k\ell} g_{\ell j} = \delta_j^k \quad . \quad (1.10)$$

We set

$$e^j = \sum_{k=1}^N g^{jk} e_k \quad , \quad e_j = \sum_{k=1}^N g_{jk} e^k \quad . \quad (1.11)$$

Then, we can expand any $x \in V$ by

$$x = \sum_{j=1}^N e_j \langle e^j | x \rangle = \sum_{j=1}^N \langle x | e_j \rangle e^j \quad . \quad (1.12)$$

We can now introduce the notion of orthogonal triple system as follows. Consider a vector space V with a triple linear product $x y z$ and with a bi-linear non-degenerate form $\langle x | y \rangle$, satisfying the axioms

$$(i) \quad \langle y | x \rangle = \langle x | y \rangle \quad (1.13a)$$

$$(ii) \quad y x z + x y z = 0 \quad (1.13b)$$

$$(iii) \quad x z y + x y z = 2\lambda \langle y | z \rangle x - \lambda \langle x | y \rangle z - \lambda \langle z | x \rangle y \quad , \quad (1.13c)$$

$$(iv) \quad \langle uvx | y \rangle = - \langle x | uvy \rangle \quad , \quad (1.13d)$$

$$(v) \quad uv(xyz) = (uvx)yz + x(uvy)z + xy(uvz) \quad , \quad (1.13e)$$

where λ is a constant. We call any such a V to be an orthogonal triple system¹.

For any orthogonal triple system, we can introduce the second triple product $[x, y, z]$ by

$$x y z = [x, y, z] + \lambda \langle y | z \rangle x - \lambda \langle z | x \rangle y \quad (1.14)$$

Then, Eqs. (1.13a)–(1.13d) imply that both $[x, y, z]$ and $\langle w | [x, y, z] \rangle$ are totally antisymmetric in x, y , and z , as well as in w, x, y , and z , respectively. However, the last relation Eq. (1.13e) will become a rather complicated equation in terms of $[x, y, z]$'s. The notion of the orthogonal triple system is a generalization of both quaternionic and octonionic triple systems. Indeed for the totally antisymmetric product $[x, y, z]$ of both quaternionic and octonionic triple systems, we conversely introduce the triple product $x y z$ by Eq. (1.14) where the constant λ is assumed to be

$$(a) \quad \lambda = \text{arbitrary for } N = 4 \quad (1.15a)$$

$$(b) \quad \lambda = -3 \beta \text{ for } N = 8 \quad . \quad (1.15b)$$

We can then verify the validity of Eqs. (1.13), so that they become orthogonal triple systems. The reason why we can choose λ to be arbitrary for the case of $N = 4$ is validities of special identities, as has been explained elsewhere¹.

2. Lie-Super Triple System

2.1 Super Space

Let the underlying vector space V admit a Z_2 -grading, so that

$$V = V_B \oplus V_F \quad (2.1)$$

is a direct sum of the bosonic vector space V_B and of the fermionic vector space V_F . We set their dimensions as

$$\text{Dim } V_B = N_B \quad , \quad \text{Dim } V_F = N_F \quad (2.2)$$

so that

$$N = N_B + N_F \quad . \quad (2.3)$$

We define the signature function $\sigma(x)$ in V by

$$\sigma(x) = \begin{cases} 0 , & \text{if } x \in V_B \\ 1 , & \text{if } x \in V_F \end{cases} \quad (2.4a)$$

$$(2.4b)$$

We consider only homogeneous elements x of V , i.e. either $x \in V_B$ or $x \in V_F$. Any triple product $x \, y \, z$ (or $[x, y, z]$) defined in V is hereafter assumed to satisfy

$$\sigma(xyz) = \{\sigma(x) + \sigma(y) + \sigma(z)\} \pmod{2} \quad . \quad (2.5)$$

Also, any bi-linear non-degenerate form $\langle x|y \rangle$ in V is called super-symmetric, if we have

$$(a) \quad \langle x|y \rangle = 0, \text{ unless } \sigma(x) = \sigma(y) \pmod{2} , \quad (2.6b)$$

$$(b) \quad \langle y|x \rangle = (-1)^{\sigma(x)\sigma(y)} \langle x|y \rangle , \quad (2.6c)$$

which we assume throughout in this note. In what follows, we often write

$$(-1)^{xy} = (-1)^{\sigma(x)\sigma(y)} \quad (2.7)$$

whenever there is no confusion.

2.2 Lie-Super Triple System

Suppose that a triple product $x \, y \, z$ satisfies the following axioms.

$$(i) \quad y \, x \, z = -(-1)^{xy} x \, y \, z, \quad (2.8a)$$

$$(ii) \quad (-1)^{xz} x \, y \, z + (-1)^{yx} y \, z \, x + (-1)^{zy} z \, x \, y = 0, \quad (2.8b)$$

$$\begin{aligned}
\text{(iii)} \quad uv(xyz) &= (uvx)yz + (-1)^{(u+v)x}x(uvy)z \\
&\quad + (-1)^{(u+v)(x+y)}xy(uvz)
\end{aligned} \tag{2.8c}$$

where we have set for simplicity

$$(-1)^{(u+v)x} = (-1)^{ux+vx} = (-1)^{[\sigma(u)+\sigma(v)]\sigma(x)}$$

etc. in accordance with Eq. (2.7). If $V = V_B$ with $V_F = 0$, then this reduces to the well-known Lie triple system^{3,4}, so that we call the system satisfying Eqs. (2.8) to be a Lie-super triple system.

The Lie-super triple system is intimately related to Lie-super algebra⁵. Let L be a Lie-super algebra with the Lie-product $[x, y]$, so that we have

$$\text{(i)} \quad \sigma([x, y]) = \{\sigma(x) + \sigma(y)\} \bmod 2 \tag{2.9a}$$

$$\text{(ii)} \quad [y, x] = -(-1)^{xy}[x, y], \tag{2.9b}$$

$$\text{(iii)} \quad (-1)^{xz}[[x, y], z] + (-1)^{yx}[[y, z], x] + (-1)^{zy}[[z, x], y] = 0. \tag{2.9c}$$

If we set

$$x \ y \ z \equiv [[x, y], z] \quad , \tag{2.10}$$

then it is not difficult to see that the triple product $x \ y \ z$ defined by Eq. (2.10) satisfies Eqs. (2.8). The converse statement is also true in a sense to be specified below.

For this, we introduce first the linear multiplication operator $L_{x,y}$ in V by

$$L_{x,y}z = x \ y \ z \quad , \tag{2.11}$$

and set

$$M = \text{vector space spanned by } L_{x,y} \text{'s} \ , \ (x, \ y \in V) \quad . \tag{2.12}$$

Calculating the commutators

$$[L_{x,y}, L_{u,v}] = L_{x,y}L_{u,v} - (-1)^{(x+y)(u+v)}L_{u,v}L_{x,y} \tag{2.13}$$

from Eq. (2.8c), we find

$$[L_{x,y}, L_{u,v}] = L_{xyu,v} + (-1)^{(x+y)u}L_{u,xyv} \tag{2.14}$$

so that this defines a Lie-super algebra with

$$\sigma(L_{x,y}) = [\sigma(x) + \sigma(y)] \bmod 2 \tag{2.15}$$

Actually, we can construct a larger Lie-super algebra in a larger space

$$V_0 = V \oplus M \quad (2.16)$$

as follows. The commutators $[M, M]$ are still defined by Eq. (2.13), while we set

$$[L_{x,y}, z] = -(-1)^{(x+y)z} [z, L_{x,y}] \equiv x \ y \ z \quad (2.17)$$

for $[M, V]$ and

$$[x, y] = -(-1)^{yx} [y, x] \equiv L_{x,y} \quad (2.18)$$

for $[V, V]$. We note especially

$$[M, M] \subset M \quad , \quad [M, V] \subset V \quad , \quad [V, V] \subset M \quad , \quad (2.19)$$

which is familiar in the theory of symmetric homogeneous spaces. It is not hard to verify that these define a Lie-super algebra in V_0 . If $V = V_B$ with $V_F = 0$, this construction reduces of course to the familiar canonical construction^{3,4}.

Then, we calculate

$$[[x, y], z] = [L_{x,y}, z] = x \ y \ z \quad (2.20)$$

from Eqs. (2.18) and (2.17), so that the relation Eq. (2.10) is formally valid, although the meaning of $[x, y]$ is different, since it is an element of M , but not of V itself.

2.3 Para-statistics as a Lie-Super Triple System

Let $\langle x_1, x_2, \dots, x_N \rangle$ be a vector space spanned by $x_1, x_2, \dots, x_N$, and set

$$V_B = \langle a_j, a_j^+ ; j = 1, 2, \dots, n \rangle \quad (2.21a)$$

$$V_F = \langle b_\alpha, b_\alpha^+ ; \alpha = 1, 2, \dots, m \rangle \quad . \quad (2.21b)$$

Moreover, we introduce $\langle x|y \rangle$ in $V = V_B \oplus V_F$ by

$$\langle a_j | a_k^+ \rangle = \langle a_k^+ | a_j \rangle = \delta_{jk} \quad , \quad (j, k = 1, 2, \dots, n) \quad , \quad (2.22a)$$

$$\langle b_\alpha | b_\beta^+ \rangle = - \langle b_\beta^+ | b_\alpha \rangle = \delta_{\alpha\beta} \quad , \quad (\alpha, \beta = 1, 2, \dots, m) \quad , \quad (2.22b)$$

while all other combinations such as $\langle a_j | a_k \rangle$, $\langle a_j | b_\alpha \rangle$, $\langle a_j | b_\alpha^+ \rangle$ etc. are assumed to be identically zero. Then, $\langle x|y \rangle$ defines a bi-linear symmetric non-degenerate form in the sense of Eqs. (2.6).

For any vector space V with $\langle x|y \rangle$ satisfying Eq. (2.6), the triple product defined by

$$x \ y \ z = \lambda \langle y|z \rangle x - \lambda(-1)^{yz} \langle x|z \rangle y \quad , \quad (2.23)$$

for a non-zero constant λ can be readily verified to give a Lie-super triple system. Applying this fact to the system $V = V_B \oplus V_F$ given by Eqs. (2.21) and (2.22), we see that V is a Lie-super triple system, while V_B itself defines a Lie triple system. Especially, for $V = V_B$, we see

$$[a_j, a_k^+, a_\ell] = [[a_j, a_k^+], a_\ell] = 2 \delta_{k\ell} a_j \quad , \quad (2.23a)$$

$$[a_j^+, a_k^+, a_\ell] = [[a_j^+, a_k^+], a_\ell] = 2 \delta_{k\ell} a_j^+ - 2 \delta_{j\ell} a_k^+ \quad , \quad (2.23b)$$

where we have set $\lambda = 2$. These relations are essentially equivalent to the para-Fermi statistics⁶. Similarly, V_F defines the para-boson statistics⁶. However, for $V = V_B \oplus V_F$, we may have rather peculiar relations such as

$$[[a_j, b_\alpha^+], a_k^+] = -2 \delta_{jk} b_\alpha^+ \quad (2.23c)$$

so that bosonic and fermionic operators are no longer independent of each other in contrast to the standard theory. This possibility will be explored, however, elsewhere. At any rate, the para-statistic relations may be rewritten as

$$[[x, y], z] = 2 \langle y|z \rangle x - 2(-1)^{yz} \langle x|z \rangle y \quad , \quad (2.24a)$$

$$[x, y] = xy - (-1)^{xy} yx \quad (2.24b)$$

where boson variables and fermion variables are now regarded to possess opposite signature values as in Eqs. (2.21) and (2.22), i.e.

$$\sigma(x) = \begin{cases} 0 & \text{for fermions} \\ 1 & \text{for bosons} \end{cases} \quad (2.25)$$

Also, this realizes a ortho-symplectic Lie-super algebra⁵.

3. Ortho-Symplectic Super-Triple System

3.1 Definition

Let $\langle x|y \rangle$ be a bi-linear non-degenerate super-symmetric form in the sense of Eqs. (2.6). Suppose now that a triple linear product $x \ y \ z$ satisfies the following axioms:

$$(i) \quad \langle y|x \rangle = (-1)^{xy} \langle x|y \rangle, \quad (3.1a)$$

$$(ii) \quad x y z + (-1)^{xy} y x z = 0, \quad (3.1b)$$

$$(iii) \quad x y z + (-1)^{yz} x z y \\ = 2 \lambda \langle y|z \rangle x - \lambda \langle x|y \rangle z - \lambda (-1)^{yz} \langle x|z \rangle y, \quad (3.1c)$$

$$(iv) \quad \langle uvx|y \rangle = -(-1)^{(u+v)x} \langle x|uvy \rangle, \quad (3.1d)$$

$$(v) \quad uv(xyz) = (uvx)yz + (-1)^{(u+v)x} x(uvy)z \\ + (-1)^{(u+v)(x+y)} xy(uvz), \quad (3.1e)$$

for a constant λ . We call the system to be ortho-symplectic super-triple system. If $V = V_B$ with $V_F = 0$, then it reduces to the orthogonal triple system, while the other case of $V = V_F$ with $V_B = 0$ is equivalent to the symplectic triple system of Yamaguchi and Asano⁷. Although the present super triple system is still a special case of more general super systems considered by many authors, it is still of some intrinsic interest as we will see below.

Here, we will construct a class of ortho-symplectic super triple system as follows. Let $\langle x|y \rangle$ be as before, and let

$$J_\alpha : V \rightarrow V \quad (\alpha = 1, 2, \dots, n) \quad (3.2)$$

be a signature-preserving linear mapping in V so that

$$\sigma(J_\alpha x) = \sigma(x) \quad . \quad (3.3)$$

Moreover, we assume the validity of

$$\langle x|J_\alpha y \rangle = - \langle J_\alpha x|y \rangle \quad (3.4)$$

as well as

$$J_\alpha J_\beta = \lambda \delta_{\alpha\beta} 1 + \sum_{\gamma=1}^n f_{\alpha\beta\gamma} J_\gamma \quad (3.5)$$

for a constant λ , where 1 stands for the identity linear mapping in V . We consider two cases of

$$(i) \quad n = 1 \quad , \quad f_{\alpha\beta\gamma} = 0, \quad (3.6a)$$

$$(ii) \quad n = 3 \quad , \quad f_{\alpha\beta\gamma} = \epsilon_{\alpha\beta\gamma} = \text{Levi-Civita symbol} \quad (3.6b)$$

so that the case of $n = 3$ corresponds to the quaternion algebra for J_α ($\alpha = 1, 2, 3$). Now, we introduce a triple product by

$$\begin{aligned} x \cdot y \cdot z = & \sum_{\alpha=1}^n \{ J_\alpha x \langle y | J_\alpha z \rangle \\ & + (-1)^{z(y+x)} J_\alpha y \langle z | J_\alpha x \rangle + J_\alpha z \langle x | J_\alpha y \rangle \} \\ & + \lambda \{ \langle y | z \rangle x - (-1)^{z(x+y)} \langle z | x \rangle y \} \quad . \end{aligned} \quad (3.7)$$

We can verify then that it satisfies Eqs. (3.1).

3.2 Its Relationship to Lie-super Triple System

Then, from a given ortho-symplectic super-triple system V , we can construct Lie-super triple systems in two different ways. The first one is essentially a straight-forward generalization of the method⁷ by Freudenthal-Yamaguchi-Asano in the space $V \oplus V$, which will be reported elsewhere. Here, we will consider a generalization of the earlier result given in ref. 8 which will be referred hereafter to as II. Let $e_1, e_2, \dots, e_N$ be a basis of V and define its dual e^j by Eq. (1.11). We introduce a new triple product in the same vector space by

$$x \cdot y \cdot z = - \sum_{j=1}^N (xye_j) e^j z - \frac{1}{3} \lambda (N_0 - 16) x \cdot y \cdot z \quad , \quad (3.8)$$

where we have set

$$N_0 = N_B - N_F \quad . \quad (3.9)$$

Then, we can show as in II that $x \cdot y \cdot z$ defines a Lie-super triple system. However, it could happen that we have $x \cdot y \cdot z = 0$ identically as in octonionic triple system as well as in other cases studied in II. As a matter of fact, the condition $x \cdot y \cdot z = 0$ is crucial to obtain some class of solutions of Yang-Baxter equation as we already noted in II.

We now calculate $x \cdot y \cdot z$ for the ortho-symplectic super-triple system given by Eq. (3.7) to find

$$x \cdot y \cdot z = c \, x * y * z \quad , \quad (3.10a)$$

$$c = \begin{cases} -\frac{1}{3} \lambda (N_0 - 4) \, , & \text{for } n = 1 \end{cases} \quad (3.10b)$$

$$c = \begin{cases} -\frac{1}{3} \lambda (N_0 + 8) \, , & \text{for } n = 3 \end{cases} \quad (3.10c)$$

where $x * y * z$ is a Lie-super triple product defined by

$$\begin{aligned}
x * y * z = \sum_{\alpha=1}^n \{ & J_{\alpha} x < y | J_{\alpha} z > \\
& + (-1)^{z(x+y)} J_{\alpha} y < z | J_{\alpha} x > - 2 J_{\alpha} z < x | J_{\alpha} y > \} \\
& + \lambda \{ < y | z > x - (-1)^{z(x+y)} < z | x > y \} \quad .
\end{aligned} \tag{3.11}$$

Especially, we see

$$x \cdot y \cdot z = 0 \tag{3.12}$$

for two cases of

$$(i) \quad n = 1 \quad , \quad N_0 = 4 \tag{3.13a}$$

$$(ii) \quad n = 3 \quad , \quad N_0 = -8 \quad . \tag{3.13b}$$

We also note that a special case of

$$x \, y \, z = \lambda \left\{ < y | z > x - (-1)^{z(x+y)} < z | x > y \right\} \tag{3.14a}$$

gives

$$x \cdot y \cdot z = -\frac{1}{3} \lambda (N_0 - 10) x \, y \, z \quad . \tag{3.14b}$$

Especially $x \, y \, z$ defined by Eq. (3.14a) describes both ortho-symplectic and Lie super triple systems at the same time.

4. Yang-Baxter Equation

Let $R_{ab}^{cd}(\theta)$ ($a, b, c, d = 1, 2, \dots, N$) be the scattering matrix for $a + b \rightarrow c + d$ in one-dimensional line with the rapidity difference θ . The (super) Yang-Baxter equation⁹ is the relation.

$$\begin{aligned}
& \sum_{a', b', c'=1}^N (-1)^{a'b' + a_2 c' + b_2 c_2} R_{a_1 b_1}^{b' a'}(\theta) R_{a' c_1}^{c' a_2}(\theta') R_{b' c'}^{c_2 b_2}(\theta'') \\
& = \sum_{a', b', c'=1}^N (-1)^{b' c' + a' c_2 + a_2 b_2} R_{b_1 c_1}^{c' b'}(\theta'') R_{a_1 c'}^{c_2 a'}(\theta') R_{a' b'}^{b_2 a_2}(\theta)
\end{aligned} \tag{4.1}$$

with the energy-momentum conservation law

$$\theta' = \theta + \theta'' \quad . \tag{4.2}$$

This relation can be graphically depicted as in Fig. 1, below.

Figure 1

Here, $(-1)^{ab} = (-1)^{\sigma(a)\sigma(b)}$ is the signature factor as before.

If we set further

$$S_{ab}^{cd}(\theta) = (-1)^{cd} R_{ab}^{cd}(\theta) \quad (4.3)$$

then we can eliminate all sign factors from Eq. (4.1) in terms of $S_{ab}^{cd}(\theta)$. However, it will lose then some interesting features of the super-space.

We assume hereafter that $R_{ab}^{cd}(\theta)$ satisfies a condition

$$R_{ab}^{cd}(\theta) = 0 \quad \text{if} \quad \sigma(c) + \sigma(d) \neq \{\sigma(a) + \sigma(b)\} \pmod{2} \quad . \quad (4.4)$$

We now introduce two θ -dependent triple linear products by

$$[e^b, e_c, e_d]_\theta = \sum_{a=1}^N e_a R_{cd}^{ab}(\theta) \quad , \quad (4.5a)$$

$$[e^a, e_d, e_c]_\theta^* = (-1)^{ab+cd} \sum_{b=1}^N e_b R_{cd}^{ab}(\theta) \quad (4.5b)$$

or equivalently by

$$\begin{aligned} R_{cd}^{ab}(\theta) &= \langle e^a | [e^b, e_c, e_d]_\theta \rangle \\ &= (-1)^{ab+cd} \langle e^b | [e^a, e_d, e_c]_\theta^* \rangle \quad . \end{aligned} \quad (4.6)$$

Note that Eqs. (4.4) and (4.5) are consistent with Eq. (2.5).

Now, the Yang-Baxter equation Eq. (4.1) can be rewritten as a triple product equation

$$\begin{aligned} & \sum_{j=1}^N (-1)^{\sigma(e_j)\sigma(z)} [v, [u, e_j, z]_{\theta'}, [e^j, x, y]_{\theta}]_{\theta''}^* \\ &= (-1)^{\sigma(u)\sigma(v)+\sigma(x)\sigma(z)} \sum_{j=1}^N (-1)^{\sigma(e_j)\sigma(x)} [u, [v, e_j, x]_{\theta'}^*, [e^j, z, y]_{\theta''}^*]_{\theta} \quad . \end{aligned} \quad (4.7)$$

Here, we will present a class of solutions of Eq. (4.7). Let $x \ y \ z$ be the ortho-symplectic super-triple product defined as in section 3. We seek a solution with ansatz of

$$\begin{aligned} [z, x, y]_{\theta} &= P(\theta)(-1)^{(x+y)z} x \ y \ z + Q(\theta) \langle x|y \rangle z \\ &\quad + R(\theta) \langle z|x \rangle y + S(\theta)(-1)^{xy} \langle z|y \rangle x \end{aligned} \quad (4.8)$$

for some functions $P(\theta)$, $Q(\theta)$, $R(\theta)$, and $S(\theta)$ of θ to be determined. We note that Eq. (4.8) implies

$$[z, x, y]_{\theta}^* = [z, x, y]_{\theta} \quad . \quad (4.9)$$

Moreover, we assume that the triple product $x \cdot y \cdot z$ defined by Eq. (3.8) vanishes identically, i.e.

$$x \cdot y \cdot z = 0 \quad . \quad (4.10)$$

Then, the solution for $P(\theta) \neq 0$ can be found to be

$$R(\theta)/P(\theta) = a + k\theta \quad , \quad (4.11a)$$

$$Q(\theta)/P(\theta) = \lambda - \frac{2a\lambda}{2(a-\lambda) + k\theta} \quad , \quad (4.11b)$$

$$S(\theta)/P(\theta) = -2\lambda - \frac{2\lambda a}{k\theta} \quad , \quad (4.11c)$$

where we have set

$$a = -\frac{1}{6} \lambda (N_0 - 4) = -\frac{1}{6} \lambda (N_B - N_F - 4) \quad , \quad (4.12)$$

and k is an arbitrary constant. Evidently, the present solution generalizes that given in II. Especially, this reproduces the solution of de Vega and Nicolai¹⁰, which is based upon the octonionic triple product corresponding to $N_B = 8$

and $N_F = 0$. Also, $x \ y \ z$ given by Eq. (3.7) with $n = 1$ satisfies $x \cdot y \cdot z = 0$ for $N_0 = 4$ with $a = 0$, so that the solution admits existence of non-zero $P(0)$, $Q(0)$, $R(0)$, and $S(0)$ for $\theta = 0$ which may be of some interest to the knot theory.

For other applications of super-triple systems, see also ref. 11.

Acknowledgements

This paper is dedicated to the 65th birthday of Professor K. C. Wali of Syracuse University. The author wishes a long productive life ahead for Professor Wali.

This paper is supported in part by the U.S. Department of Energy Grant No. DE-FG-02-91ER40685.

References

1. S. Okubo, *Jour. Math. Phys.* in press.
2. R. D. Schafer, *An Introduction to Non-associative Algebras* (Academic, New York, 1966).
3. W. G. Lister, *Am. J. Math.* **89** (1952), 787.
4. K. Yamaguchi, *J. Sci. Hiroshima University* **A21**, (1958) 155.
5. M. Scheunert, *The Theory of Lie Superalgebra* (Springer-Verlag, Berlin-Heidelberg-New York, 1979).
6. Y. Ohnuki and S. Kamefuchi, *Quantum Field Theory and Parastatistics* (University of Tokyo Press, Tokyo/Springer-Verlag, Berlin, 1982).
7. K. Yamaguchi and H. Asano, *Proc. Jap. Acad.* **51**, (1972) 247.
8. S. Okubo, *Jour. Math. Phys.* in press.
9. *Yang-Baxter Equation in Integrable Systems*, edited by M. Jimbo (World Scientific, Singapore, 1989).
10. H. J. de Vega and H. Nicolai, *Phys. Lett.* **B244**, (1990) 295.
11. M. Günaydin, *Phys. Lett.* **B255**, (1991) 46. M. Günaydin and S. Hyun, *Mod. Phys. Lett.* **6**, (1991) 1733 and *Nucl. Phys.* **B373**, (1992) 688.